## What is HIV?

HIV (Human Immunodeficiency Virus) is a virus that attacks the immune system, weakening the body's ability to fight infections and diseases. Without proper treatment, HIV can progress to AIDS (Acquired Immunodeficiency Syndrome).

#### **Key Facts**

- Causes: HIV is caused by a virus that attacks the body's immune system, specifically targeting and destroying CD4 cells, also known as T-helper cells.
- Transmission: HIV is primarily spread through unprotected sex, sharing needles, blood transfusion and from mother to child during childbirth or breastfeeding. HIV is not spread through casual contact, such as hugging, shaking hands, or sharing utensils.
- Effects on the Immune System: HIV targets CD4 cells (T cells), leading to a compromised immune system that cannot defend against opportunistic infections.




## What is Hypertension?

Hypertension, or high blood pressure, is a condition where the force of blood against the artery walls is too high.

Often called the "silent killer," it can lead to serious health problems if untreated.

## **Key Facts**

Risk Factors: Poor diet (high in salt and fat), lack of exercise, smoking, excessive alcohol consumption, stress, and genetic predisposition.



Health Implications:
Uncontrolled hypertension
increases the risk of heart
disease, stroke, kidney damage,
and other complications.





#### **Understanding**

HIV and Hypertension

# Protecting Your Health,

**One Step at a Time** 





# Your Health, Your Responsibility

- Get tested regularly for HIV and monitor your blood pressure.
- Adopt a heart-healthy lifestyle: Eat balanced meals, stay active, and avoid smoking or excessive alcohol.
- Consult your healthcare provider about managing both conditions effectively.

**Stay informed. Take** control. Live healthy.

## Why It Matters?

Managing both conditions together is critical for longterm health. Early diagnosis, regular monitoring, and lifestyle changes can significantly improve outcomes.







## **How Are HIV and Hypertension Connected?**

People living with HIV are at a higher risk of developing hypertension due to factors such as:



#### **Medication Side Effects**

Some antiretroviral therapies (ART) can increase blood pressure.

#### **Chronic Inflammation**



HIV causes inflammation, which can damage blood vessels over time.

#### **Aging with HIV**



Aging increases hypertension risk, especially with HIV and ART effects The significantly increases hypertension risk

## **Lifestyle Factors**



Smoking, poor diet, and inactivity can increase risks for both conditions.

#### **Co-infection and Comorbidities**



Co-infections like Hepatitis C and Chronic Light Midney Disease (CKD) are common in HIV patients. These co-infections worsen vascular damage, increasing HIV-related hypertension.